

539,415

10/539415

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
8 July 2004 (08.07.2004)

PCT

(10) International Publication Number
WO 2004/056795 A1

(51) International Patent Classification⁷: **C07D 333/20**,
A61K 31/381, A61P 25/24

A.A. Thornton & Co., 235 High Holborn, London WC1V 7LE (GB).

(21) International Application Number:
PCT/GB2003/005357

(74) Agents: WAIN, Christopher, Paul et al.; A.A. Thornton & Co., 325 High Holborn, London WC1V 7LE (GB).

(22) International Filing Date:
10 December 2003 (10.12.2003)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0229583.0 19 December 2002 (19.12.2002) GB

(84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): CIPLA LTD [IN/IN]; 289 Bellasis Road, Mumbai Central, Mumbai 400 008 (IN).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): RAO, Dharmaraj, Ramachandra [IN/IN]; 4/403 Garden Enclave, Pokhran Road 2, Thane (West), Mumbai 400 601, Maharashtra (IN). KANKAN, Rajendra, Narayanrao [IN/IN]; A-3/5, NBD Society, NSS Road, Gratkopar (West), Mumbai 400084, Maharashtra (IN). WAIN, Christopher, Paul [GB/GB];

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: A PROCESS FOR PREPARING DULOXETINE AND INTERMEDIATES FOR USE THEREIN

(57) Abstract: A process for preparing (+)duloxetine, or an acid addition salt thereof, which process comprises resolving racemic (±)duloxetine with a chiral acid so as to obtain a salt of the chiral acid and (+)duloxetine, substantially free of (-)duloxetine; and (ii) if desired, converting the salt prepared in step (i) to the free base or another acid addition salt as appropriate. The process for preparing (+)duloxetine, or an acid addition salt thereof, can further comprise an O-alkylation intermediate process step which is carried out in the presence of a base and a phase transfer catalyst.

WO 2004/056795 A1


A PROCESS FOR PREPARING DULOXETINE AND
INTERMEDIATES FOR USE THEREIN

The present invention is concerned with a process for preparing duloxetine, in particular a process for preparing (+)duloxetine in enantiomerically pure form, and to intermediates for use therein.

Duloxetine, N-methyl-3-(1-naphthalenyloxy)-3-(2-thienyl)propanamine, is a dual serotonin and norepinephrine reuptake inhibitor. (+)Duloxetine has particular therapeutic utility as an anti-depressant.

Duloxetine, and the preparation thereof, is described in US patents 5023269 and 4956388, and also Tetrahedron Letters, 31, (49), 7101-04, 1990. Seven different routes of synthesis have also been reported in Drugs of the Future 2000, 25(9) 907-916. These syntheses have involved either a resolution of a key intermediate or a stereospecific reduction of a keto group to the alcohol.

We have now observed that a particular chiral intermediate employed in the synthesis of (+)duloxetine, namely



undergoes considerable racemisation during coupling with a 1-naphthyl halide under the reaction conditions employed in the prior art. In particular, in the presence of a strong base, such as sodium hydride, and a protic polar solvent, such as DMSO, the dimethyl anion can be generated which can cause partial or complete racemisation of the condensed product.

There is, therefore, a need for an improved process for the preparation of (+)duloxetine which alleviates the problems associated with prior art processes as referred to above. We have developed such a process which is advantageous in obviating racemisation, so as to yield an enantiomerically pure form of (+)duloxetine. In particular, our process can be seen to achieve the above described advantage, by carrying out resolution as a final step in the reaction process (thereby obviating the opportunity for racemisation during preceeding intermediate process steps) and / or avoiding conditions that would result in the production of intermediate products that would be prone to racemisation.

According to the present invention, therefore, there is provided a process for preparing (+)duloxetine, or an acid addition salt thereof, which process comprises:

- (i) resolving racemic (\pm)duloxetine with a chiral acid so as to obtain a salt of the chiral acid and (+)duloxetine, substantially free of (-)duloxetine; and
- (ii) if desired, converting the salt prepared in step (i) to the free base or another acid addition salt as appropriate.

The resolution step (i) is achieved with a suitable chiral acid in a suitable solvent. The chiral acid can typically be selected from the group consisting of mandelic acid, tartaric acid, di-p-toluyyl tartaric acid, dibenzoyl tartaric acid, camphor sulfonic acid and the like. Other suitable chiral acids may be determined by testing and the use thereof in a process as described above falls within the scope of the present invention. Preferably the chiral acid employed in a process according to the present invention is (-)di-p-toluyyl tartaric acid. Suitably, the solvent employed is a lower alkanol, such as methanol or ethanol, although again other suitable solvents can be determined by testing and the use thereof in a process as described above falls within the scope of the present invention. A preferred solvent is methanol.

The salts of (+)duloxetine prepared by resolution step (i) represent a further aspect of the present invention and there is further provided by the present invention, therefore, a salt of a chiral acid and (+)duloxetine, substantially free of (-)duloxetine. Such salts of a chiral acid

and (+)duloxetine, substantially free of (-)duloxetine, are useful as intermediates for preparing the free base or another acid addition salt as appropriate.

Suitable salts provided by the present invention include (+)duloxetine mandelate, (+)duloxetine tartrate, (+)duloxetine di-p-toluy l tartrate, (+)duloxetine dibenzoyl tartrate, (+)duloxetine camphor sulfonate and the like. A preferred salt according to the present invention is (+)duloxetine di-p-toluy l tartrate, which is useful as an intermediate for preparing the free base or another acid addition salt as appropriate.

Intermediate salts prepared according to the present invention as described above can be converted to the free base or another acid addition salt according to step (ii) of a process according to the present invention. Suitably, an intermediate salt of the chiral acid and (+)duloxetine can be treated with a base, such as sodium hydroxide, to yield the free base. The free base itself can, if desired, be converted into an acid addition salt thereof.

Suitable acid addition salts which may be formed in step (ii) include those formed with pharmaceutically acceptable organic or inorganic acids and are well known to those of skill in the art. Acids commonly employed to form such salts include inorganic acids such as hydrochloric, hydrobromic, hydroiodic, sulfuric and phosphoric acid, as well as organic acids such as para-toluenesulfonic, methanesulfonic, oxalic, para-bromophenylsulfonic, carbonic, succinic, citric, benzoic and acetic acid, and related inorganic and organic acids. Such pharmaceutically acceptable salts thus include sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caprate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, terephthalate, sulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, β -hydroxybutyrate, glycollate, maleate, tartrate, methanesulfonate, propanesulfonates, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate and the like salts. Preferred pharmaceutically acceptable acid addition salts include those formed with mineral acids such as hydrochloric acid and hydrobromic acid, and those formed with organic

acids such as oxalic acid and maleic acid. A particularly preferred acid addition salt is the hydrochloride.

A particularly preferred process according to the present invention comprises:

- (i) resolving racemic (\pm)duloxetine with di-p-toluyyl tartaric acid so as to obtain (+)duloxetine di-p-toluyyl tartrate, substantially free of (-)duloxetine; and
- (ii) converting (+)duloxetine di-p-toluyyl tartrate prepared in step (i) to (+)duloxetine hydrochloride.

A process according to the present invention preferably yields (+)duloxetine in substantially pure enantiomeric form. Thus the ratio of (+)duloxetine: (-)duloxetine as prepared by the present invention may be at least about 94:6, such as at least about 98:2, or more preferably at least about 99:1. Preferably (+)duloxetine prepared by a process according to the present invention has an enantiomeric purity of at least about 99%, or more particularly at least about 99.5%.

The proportion of (+)duloxetine achieved by a process according to the present invention can be increased by effecting one or more, for example two or three, recrystallisations in step (i). To this end, a crystalline salt obtained as a result of resolution may be dissolved in a solvent therefor and the salt recrystallised from the resulting solution. In this way the proportion of a salt with (+)duloxetine can thus be increased until it is substantially pure, that is substantially only a salt with (+)duloxetine is present.

The mother liquor from resolution step (i), or the mother liquor from the or each recrystallisation step, is enriched with (-)duloxetine. (-)Duloxetine present in one or more of these liquors, or the pooled liquors, may be converted into (\pm)duloxetine for reuse in a process according to the present invention substantially as hereinbefore described.

A further preferred aspect of a process according to the present invention comprises:


(i) resolving racemic (\pm)duloxetine with a chiral acid and obtaining a mother liquor enriched in (-)duloxetine;

(ii) converting (-)duloxetine obtained from step (i) to (\pm)duloxetine; and

(iii) if desired, employing (\pm)duloxetine obtained from step (ii) in a process according to the present invention substantially as hereinbefore described.

Suitably, one or more mother liquors obtained from a process as described above, or pooled such mother liquors, may be treated with a base to remove any residual chiral acid and to thereby afford the free base enriched in (-)duloxetine. The free base can then be converted to the racemate, typically by reflux in a suitable solvent for several hours, optionally in the presence of a suitable acid (e.g. HCl) or a base (e.g. NaOH), which racemate can then be recycled for use in a process according to the present invention substantially as hereinbefore described.


Substantially as hereinbefore described an aim of a process according to the present invention is to obviate racemisation associated with prior art processes so as to yield an enantiomerically pure form of (+)duloxetine. According to the present invention this can be achieved by the resolution of (\pm)duloxetine as a final process step in the preparation of (+)duloxetine substantially as hereinbefore described, and according to a preferred aspect of the present invention the prior art techniques for the preparation of duloxetine can also be further modified by use of the following reaction conditions in an O-alkylation step employed in the synthesis of duloxetine. More particularly, an intermediate process step employed in a process according to the present invention comprises reacting intermediate compounds of formulae (I) and (II) so as to yield a compound of formula (III), or an acid addition salt thereof:



in the presence of a base and a phase transfer catalyst, where one of X and Y is hydroxy and the other is a leaving group. The use of such reaction conditions according to the present invention is advantageous in avoiding the use of stronger bases, such as sodium hydride, employed according to prior art preparatory techniques (such stronger bases being a fire hazard and also expensive), with the use of phase transfer catalysts facilitating the above reaction step to be carried out under milder conditions.

The use of the above reaction conditions can also offer advantages when chiral intermediates are employed in the synthesis of (+)duloxetine, by avoiding the considerable racemisation of the chiral intermediate (S)-2-(1-N,N-dimethyl-3-hydroxy)propyl amino)-thiophene seen

during coupling thereof with a 1-naphthyl halide under the reaction conditions employed in the prior art. In particular, as previously discussed in the presence of a strong base, such as sodium hydride, and a protic polar solvent, such as DMSO, the dimesyl anion was generated under the prior art reaction conditions, which caused partial or complete racemisation of the condensed product. This undesirable racemisation can be avoided by carrying out the O-alkylation step under the reaction conditions of the present invention and there is, therefore, further provided a process for preparing (+)duloxetine, or an acid addition salt thereof, which comprises the following intermediate process step of reacting intermediate compounds of formulae (Ia) and (II) so as to yield a compound of formula (IIIa), or an acid addition salt thereof:



Preferably the base employed can be an alkali metal hydroxide, an alkali metal carbonate, an alkali metal bicarbonate or the like. Suitably the base can be selected from the group consisting of potassium hydroxide, sodium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate and the like. Potassium hydroxide can often be preferred.

Suitable phase transfer catalysts can be crown ethers, quaternary ammonium salts, quaternary phosphonium salts and the like. A preferred phase transfer catalyst can suitably be tetrabutyl ammonium bromide or 18-Crown-6.

A suitable leaving group as represented by X or Y can be p-toluenesulfonyl, methanesulfonyl, triphenylphosphine oxide, halo and the like, with halo being preferred.

Preferably X is hydroxy and Y is a leaving group such as halo, typically fluoro.

An intermediate of formula (III), or an acid addition salt thereof, is typically converted to (\pm)duloxetine by demethylating, typically in the presence of a reagent such as a phenyl chloroformate or trichloroethyl chloroformate or the like to provide the corresponding intermediate, which is then hydrolysed to provide (\pm)duloxetine, or an acid addition salt thereof. An intermediate of formula (IIIa), or an acid addition salt thereof, is similarly typically converted to (+)duloxetine, or an acid addition salt thereof, by demethylating, typically in the presence of a reagent such as a phenyl chloroformate or trichloroethyl chloroformate or the like to provide the corresponding intermediate, which is then hydrolysed to provide (+)duloxetine, or an acid addition salt thereof.

A further preferred process according to the present invention comprises reacting intermediate compounds of formulae (I) and (II) to yield an intermediate compound of formula (III), or an acid addition salt thereof, demethylating a compound of formula (III), or an acid addition salt, so as to yield (\pm)duloxetine and converting (\pm)duloxetine to (+)duloxetine, or an acid addition salt thereof, employing a process according to the present invention substantially as hereinbefore described. A still further preferred process according to the present invention comprises reacting intermediate compounds of formulae (Ia) and (II) to yield an intermediate compound of formula (IIIa), or an acid addition salt thereof,

demethylating a compound of formula (IIIa), or an acid addition salt, so as to yield (+)duloxetine, or an acid addition salt thereof.

A particularly preferred process according to the present invention comprises reacting intermediate compounds of formulae (I) and (II) where X is hydroxy and Y is fluoro, followed by oxalic acid to yield an oxalate salt of intermediate compound of formula (III), demethylating the intermediate compound of formula (III) so as to yield (±)duloxetine and converting (±)duloxetine to (+)duloxetine employing a process according to the present invention substantially as hereinbefore described. In particular it is especially preferred that resolution to yield (+)duloxetine comprises:

- (i) resolving racemic (±)duloxetine with (-)di-p-toluyll tartaric acid so as to obtain (+)duloxetine di-p-toluyll tartrate, substantially free of (-)duloxetine; and
- (ii) converting (+)duloxetine di-p-toluyll tartrate prepared in step (i) to (+)duloxetine hydrochloride.

The compounds employed as starting materials in the processes of the present invention can be prepared by standard procedures known in the art.

The present invention further provides (+)duloxetine, or an acid addition salt thereof, obtained by a process substantially as hereinbefore described. Such (+)duloxetine, or an acid addition salt thereof, according to the present invention can suitably be formulated to provide a pharmaceutical composition according to the present invention, and there is further provided by the present invention a pharmaceutical composition comprising (+)duloxetine, or an acid addition salt thereof, obtained by a process substantially as hereinbefore described.

The present invention will now be illustrated by the following Examples, which do not limit the scope of the invention in any way.

Example 1

N,N-dimethyl-?-(1-naphthalenyloxy)-2-thiophenepropanamine oxalate

2-(1-N,N-dimethyl-3-hydroxy)propyl amino)-thiophene (12.4gms) was dissolved in DMSO (70ml). Potassium hydroxide (18.7gms) and tetrabutyl ammonium bromide were added (0.1gms). The reaction mixture was stirred at 60EC for 1 hour and 4-fluoro naphthalene (11.7gms) was added slowly over 2 hours. After the reaction was complete, it was quenched in ice-water and extracted into toluene. The toluene layer was dried and concentrated under vacuum. The residue was dissolved in 100ml of ethyl acetate and oxalic acid (7gms) was added. The reaction mixture was stirred for 1 hour and filtered (15.5gms, 76% yield).

Example 2

(±)-N-methyl-?-(1-naphthalenyloxy)-2-thiophenepropanamine ((±) Duloxetine)

N,N-dimethyl-?-(1-naphthalenyloxy)-2-thiophenepropanamine (50gms) was dissolved in toluene (250ml). To the reaction mixture was added diisopropyl ethyl amine (24.8gms) and phenyl chloroformate (30gms) was added slowly. The reaction mixture was heated to 60EC and stirred for 2 hours. After completion of the reaction, the mixture was quenched in 50ml of 5% sodium bicarbonate solution. The toluene layer was separated, dried and concentrated to give a residue.

The residue was dissolved in a mixture of DMSO and water. To this was added sodium hydroxide (22gms) and the reaction mixture refluxed for 8 hours. The reaction was then diluted with 700ml of water and extracted with ethyl acetate. The ethyl acetate layer was dried and concentrated to give the title compound (74%).

Example 3

(+)-N-methyl-?-(1-naphthalenyloxy)-2-thiophenepropanamine ((+) Duloxetine)

(±) Duloxetine (20gms) was dissolved in methanol (20ml) and (-)-di-p-toluyyl tartaric acid (6.5gms) was added. The reaction was stirred under reflux for 1 hour, methanol concentrated under vacuum, acetone (150ml) charged and cooled to 5EC. The solids obtained were filtered and dried under vacuum at 40EC to give the (-)-di-p-toluyyl tartrate salt of (+) duloxetine (12.5 g).

The above tartrate salt was suspended in a mixture of water and toluene and 500mg of sodium hydroxide was added under stirring. The toluene layer was separated, dried and concentrated under vacuum to residue. Ethyl acetate (20ml) was added to the residue followed by oxalic acid (2.2gms). The reaction was cooled to 5EC and the mixture stirred for 1 hour. The solids were filtered and dried under vacuum at 40EC to yield the title compound as an oxalate (6.5gms).

The methanolic mother liquor obtained after filtration of the (-)DPTA salt of (+)duloxetine was basified to pH 10 by lye solution (50% NaOH) and heated to reflux for 10 hours. Extractions with dichloromethane (100 ml) followed by concentration of the solvent yielded the racemate base (8.5 gm) having enantiomeric ratio (+/-) of 48:52 by chiral HPLC.

Example 4

(S)-N,N-dimethyl-1-(1-naphthalenyloxy)-2-thiophenepropanamine oxalate

(S)-2-(1-N,N-dimethyl-3-hydroxy)propyl amino)-thiophene (12.5gms) was dissolved in DMSO (60ml). Potassium hydroxide (20.5 gms) and 18-Crown-6 were added (0.12gms). The reaction mixture was stirred at 60°C for 1 hour and 4-fluoro naphthalene (11.9gms) was added slowly over 2 hours. After the reaction was complete, it was quenched in ice-water and extracted into toluene. The toluene layer was dried and concentrated under vacuum. The residue was dissolved in 100ml of ethyl acetate and oxalic acid (7.5 gms) was added. The reaction mixture was stirred for 1 hour and filtered to give the title compound in 88% yield.

Example 5

(+)-N-methyl-1-(1-naphthalenyloxy)-2-thiophenepropanamine hydrochloride ((+) Duloxetine hydrochloride)

(S)-N,N-dimethyl-1-(1-naphthalenyloxy)-2-thiophenepropanamine oxalate (50gms) was dissolved in water. The pH of the solution was adjusted to between 9 –10 with 5% sodium hydroxide. The mixture was extracted with toluene. The toluene layer was concentrated under

vacuum to about 250 ml. To the reaction mixture was added ? -chloroethyl chloroformate (30gms) slowly. The reaction mixture was heated to 60°C and stirred for 2 hours. After completion of the reaction, the mixture was concentrated under vacuum to a residue.

The residue was dissolved in methanol and refluxed for 2 hours. The reaction mixture was concentrated under vacuum to give a solid residue which was triturated with acetone to give the title compound (36 gms).

CLAIMS:


1. A process for preparing (+)duloxetine, or an acid addition salt thereof, which process comprises:
 - (i) resolving racemic (\pm)duloxetine with a chiral acid so as to obtain a salt of the chiral acid and (+)duloxetine, substantially free of (-)duloxetine; and
 - (ii) if desired, converting the salt prepared in step (i) to the free base or a further acid addition salt.
2. A process according to claim 1, wherein the chiral acid is selected from the group consisting of mandelic acid, tartaric acid, di-p-toluyyl tartaric acid, dibenzoyl tartaric acid and camphor sulfonic acid.
3. A process according to claim 2, wherein the chiral acid is di-p-toluyyl tartaric acid.
4. A process according to any of claims 1 to 3, wherein step (ii) comprises reacting a salt prepared in (i) with hydrochloric acid to yield (+)duloxetine hydrochloride.
5. A process for preparing (+)duloxetine hydrochloride, which process comprises:
 - (i) resolving racemic (\pm)duloxetine with di-p-toluyyl tartaric acid so as to obtain (+)duloxetine di-p-toluyyl tartrate, substantially free of (-)duloxetine; and
 - (ii) converting (+)duloxetine di-p-toluyyl tartrate prepared in step (i) to (+)duloxetine hydrochloride.
6. A process which comprises:

(i) resolving racemic (\pm)duloxetine with a chiral acid in a process according to any of claims 1 to 5, and obtaining a mother liquor enriched in (-)duloxetine;

(ii) converting (-)duloxetine obtained from step (i) to (\pm)duloxetine; and


(iii) if desired, employing (\pm)duloxetine obtained from step (ii) in a process according to any of claims 1 to 5.

7. A process for making (+)duloxetine or an acid addition salt thereof, comprising reacting intermediate compounds of formulae (I) and (II) so as to yield a compound of formula (III), or an acid addition salt thereof:



in the presence of a base and a phase transfer catalyst, where one of X and Y is hydroxy and the other is a leaving group, and subjecting a compound of formula (III), or an acid addition salt thereof, to further process steps so as to yield (+) duloxetine, or an acid addition salt thereof, substantially free of (-)duloxetine.

8. A process for making (+)duloxetine or an acid addition salt thereof, comprising reacting intermediate compounds of formulae (Ia) and (II) so as to yield a compound of formula (IIIa), or an acid addition salt thereof:




in the presence of a base and a phase transfer catalyst, where one of X and Y is hydroxy and the other is a leaving group, and subjecting a compound of formula (IIIa), or an acid addition

salt thereof, to further process steps so as to yield (+) duloxetine, or an acid addition salt thereof, substantially free of (-)duloxetine.

9. A process according to claim 7 or 8, wherein an intermediate of formula (III) or (IIIa), or an acid addition salt thereof, is converted to (\pm)duloxetine, or (+)duloxetine respectively, by demethylating.

10. A process for making (+)duloxetine or an acid addition salt thereof, which comprises:


(i) reacting intermediate compounds of formulae (I) and (II) to yield an intermediate compound of formula (III), or an acid addition salt thereof,



in the presence of a base and a phase transfer catalyst;

- (ii) demethylating a compound of formula (III), or an acid addition salt, so as to yield (\pm)duloxetine; and
- (iii) converting (\pm)duloxetine obtained in step (ii) to (+)duloxetine, or an acid addition salt thereof, employing a process according to any of claims 1 to 5.
11. A process for making (+)duloxetine hydrochloride which comprises:

- (i) reacting intermediate compounds of formulae (I) and (II), where X is hydroxy and Y is fluoro, followed by oxalic acid to yield an oxalate salt of intermediate compound of formula (III)



in the presence of a base and a phase transfer catalyst;

(ii) demethylating said intermediate compound of formula (III) obtained by step (i) so as to yield (\pm)duloxetine; and

(iii) converting (\pm)duloxetine obtained in step (ii) to (+)duloxetine by resolving racemic (\pm)duloxetine with di-p-toluyyl tartaric acid so as to obtain (+)duloxetine di-p-toluyyl tartrate, substantially free of (-)duloxetine, and converting said (+)duloxetine di-p-toluyyl tartrate to (+)duloxetine hydrochloride.

12. A process according to any of claims 7 to 11, wherein the base is selected from the group consisting of an alkali metal hydroxide, an alkali metal carbonate and an alkali metal bicarbonate.

13. A process according to claim 12, wherein the base is selected from the group consisting of potassium hydroxide, sodium hydroxide, potassium carbonate, sodium carbonate and sodium bicarbonate.

14. A process according to any of claims 7 to 13, where the phase transfer catalyst is selected from the group consisting of crown ethers, quaternary ammonium salts and phosphonium salts.

15. A process according to claims 7 to 10, wherein X is hydroxy and Y is a leaving group.

16. A process according to claim 15, wherein the leaving group is halo.

17. A process according to claim 16, wherein the leaving group is fluoro.

18. A salt of a chiral acid and (+)duloxetine, substantially free of (-)duloxetine.

19. A salt of a chiral acid and (+)duloxetine, substantially free of (-)duloxetine, selected from the group consisting of (+)duloxetine mandelate, (+)duloxetine tartrate, (+)duloxetine di-p-toluyyl tartrate, (+)duloxetine dibenzoyl tartrate and (+)duloxetine camphor sulfonate.

20. (+)duloxetine di-p-toluyyl tartrate, substantially free of (-)duloxetine.
21. (+)duloxetine, or an acid addition salt thereof, obtained by a process according to any of claims 1 to 17.
22. A pharmaceutical composition comprising (+)duloxetine, or an acid addition salt thereof, obtained by a process according to any of claims 1 to 17.

INTERNATIONAL SEARCH REPORT

Application No

PCT/GB 03/05357

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D333/20 A61K31/381 A61P25/24

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 457 559 A (LILLY CO ELI) 21 November 1991 (1991-11-21) page 3, line 48 -page 5; claims; example 1 ---	1,2, 4-18,22
Y	WO 00 61540 A (LILLY CO ELI ;KJELL DOUGLAS PATTON (US); LORENZ KURT THOMAS (US)) 19 October 2000 (2000-10-19) page 6, line 24 - page 8; page 11, lines 12-22; claims ---	1,2, 4-18,22
A	US 5 362 886 A (BERGLUND RICHARD A) 8 November 1994 (1994-11-08) column 1, line 48-56 -column 2; claims; examples 1,2 --- -/--	1-22

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

2 April 2004

Date of mailing of the international search report

15/04/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Chouly, J

INTERNATIONAL SEARCH REPORT

Int. Patent Application No
PCT/GB 03/05357

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 273 658 A (LILLY CO ELI) 6 July 1988 (1988-07-06) claims; examples 1,2,8,9,14 & US 5 023 269 A 11 June 1991 (1991-06-11) cited in the application ---	1-22
A	DEETER J ET AL: "ASYMMETRIC SYNTHESIS AND ABSOLUTE STEREOCHEMISTRY OF LY248686" TETRAHEDRON LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 31, no. 49, 26 November 1990 (1990-11-26), pages 7101-7104, XP001119089 ISSN: 0040-4039 cited in the application the whole document ---	1-22
P,Y	WO 03 062219 A (BORGHESE ALFIO ;LILLY CO ELI (US)) 31 July 2003 (2003-07-31) claims; examples -----	1,2, 4-18,22

INTERNATIONAL SEARCH REPORT

Application No
PCT/GB 03/05357

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0457559	A	21-11-1991	CA 2042346 A1 EP 0457559 A2 FI 912280 A HU 57760 A2 IE 911677 A1 JP 4226948 A	18-11-1991 21-11-1991 18-11-1991 30-12-1991 20-11-1991 17-08-1992
WO 0061540	A	19-10-2000	AU 3877500 A CA 2362185 A1 EP 1171417 A1 JP 2002541235 T WO 0061540 A1 US 6541668 B1	14-11-2000 19-10-2000 16-01-2002 03-12-2002 19-10-2000 01-04-2003
US 5362886	A	08-11-1994	AT 199084 T AU 685494 B2 AU 7572094 A BR 9404045 A CA 2133899 A1 CN 1109470 A ,B CZ 9402465 A3 DE 69426663 D1 DE 69426663 T2 DK 650965 T3 EP 0650965 A1 ES 2153850 T3 FI 944773 A GR 3035715 T3 HU 68943 A2 IL 111188 A JP 7188065 A NO 943825 A NZ 264633 A PL 305326 A1 PT 650965 T RU 2127269 C1 SI 650965 T1 TW 381090 B US 5491243 A ZA 9407839 A	15-02-2001 22-01-1998 04-05-1995 13-06-1995 13-04-1995 04-10-1995 17-05-1995 15-03-2001 21-06-2001 26-02-2001 03-05-1995 16-03-2001 13-04-1995 31-07-2001 28-08-1995 15-06-1998 25-07-1995 18-04-1995 22-09-1997 18-04-1995 31-05-2001 10-03-1999 30-06-2001 01-02-2000 13-02-1996 09-04-1996
EP 0273658	A	06-07-1988	AT 57924 T AU 591007 B2 AU 8266087 A CA 1302421 C CN 87108175 A ,B CY 1682 A DE 3765919 D1 DK 664887 A EG 18230 A EP 0273658 A1 GR 3001207 T3 HK 69693 A HU 47561 A2 IL 84863 A JP 2549681 B2 JP 63185946 A KR 9603808 B1	15-11-1990 23-11-1989 23-06-1988 02-06-1992 06-07-1988 10-10-1993 06-12-1990 23-06-1988 30-10-1992 06-07-1988 30-07-1992 30-07-1993 28-03-1989 29-03-1992 30-10-1996 01-08-1988 22-03-1996

INTERNATIONAL SEARCH REPORT

Int. ... onal Application No
PCT/GB 03/05357

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0273658	A	MX 9845 A	01-12-1993
		NZ 222980 A	28-11-1989
		PH 26556 A	19-08-1992
		PT 86389 A , B	01-01-1988
		SG 114992 G	29-01-1993
		SU 1598865 A3	07-10-1990
		US 4956388 A	11-09-1990
		US 5023269 A	11-06-1991
		ZA 8709472 A	30-08-1989
WO 03062219	A	31-07-2003	WO 03062219 A1
			31-07-2003